

# **Egyptian Fertility Sterility Society**

## **The 26<sup>th</sup> Editorial**

### **Does antenatal corticosteroid reduce the risk of neonatal respiratory distress syndrome in late preterm birth twin neonates?**

**Prof. Ibrahim Mahrous, Al Azhar University**

**Dr. Mohamed El Sherbiny, MOH**

#### **What Is Known Already?**

In singleton pregnancies at risk of early preterm delivery, antenatal corticosteroid (ANC) administration decreases the risk of neonatal mortality and morbidities, such as respiratory distress syndrome (RDS).<sup>6</sup> [1] Therefore, routine administration of ANC to singleton pregnant women at risk of early preterm delivery (24 to 34 weeks' gestational age) within 7 days is recommended. [1]. In contrast the efficacy of ANC administration in twin pregnancies are few and mostly performed retrospectively with conflicting findings. [2-4]

#### **What Is New?**

A recent multicenter randomized trial, twin-pregnant women at 34 weeks 0 days to 36 weeks 5 days of gestation at risk of late preterm delivery were enrolled across 8 university-based clinical centers [5]. A total of 812 participants were randomized and analyzed, with 410 receiving antenatal betamethasone in the intervention group and 402 in the placebo group. Among 1620 neonates (818 in the intervention group and 802 in the placebo group), there were no perinatal

## Conclusions.

In this randomized control trial, ANC in women with twin pregnancies at risk of late preterm delivery significantly reduced the risk of neonatal RDS. The outcomes from this study could serve as a valuable reference in counselling clinical management of twin pregnancies at risk of late preterm delivery.

..

## References

- 1- McGoldrick E ,McGoldrick E, Stewart F, Parker R, Dalziel SR. . Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev.* (12):2020. ,
- 2- Melamed N, Shah J, Yoon EW, et al. The role of antenatal corticosteroids in twin pregnancies complicated by preterm birth. *Am J Obstet Gynecol.* Lee SM, Park HS, Choi SR, Lee J, Kim HJ, Park JY, Oh KJ, Cho GJ, Oh MJ, Chung JH, Kim SM, Kim BJ, Kim SY, Hong S, Jung YM, Lee SJ, Seong JS, Kim H, Oh S, Lee J, Jin YR, Kim JH, Cho HY, Park CW, Park JS, Jun 2016;215(4):482e1-482e9. doi:10.1016/j.ajog.2016.05.037
- 3- Assad Gonçalves-FerriW, Martinez FE, Martins-Celini FP, et al. Evaluation of the effectiveness of antenatal corticoid in preterm twin and single pregnancies: a multicenter cohort study. *JMatern Fetal Neonatal Med.*;35(18): 2022 3502-3508. doi:10.1080/14767058.2020.1822806
- 4- Ushida T, Kotani T, Sadachi R, et al; Neonatal Research Network of Japan. Antenatal corticosteroids and outcomes in preterm twins. *Obstet. Gynecol.* 135(6):1387-1397. 2020
- 5- Lee SM, Park HS, Choi SR, Lee J, Kim HJ, Park JY, Oh KJ, Cho GJ, Oh MJ, Chung JH, Kim SM, Kim BJ, Kim SY, Hong S, Jung YM, Lee SJ, Seong JS, Kim H, Oh S, Lee J, Jin YR, Kim JH, Cho HY, Park CW, Park JS, Jun Antenatal Corticosteroid in Twin-Pregnant Women at Risk of Late Preterm Delivery: A Randomized Clinical Trial. *JAMA Pediatr.*;179(12):1275. 2025